

## Studies on different levels and sources of Nitrogen in augmenting the growth, yield and economics of fodder maize (*Zea mays* L.)

\*S. N. Srikantha<sup>1</sup>, S. Natarajan<sup>1</sup>, P. Anandan<sup>1</sup> and K. Arivazhagan<sup>2</sup>

<sup>1</sup>Department of Agronomy,

<sup>2</sup>Department of Soil Science & Agricultural Chemistry, Faculty of Agriculture, Annamalai University Annamalai Nagar - 608002 (India)

\*Correspondence author e-mail: [srikanthsanju07@gmail.com](mailto:srikanthsanju07@gmail.com)

contact: 6363628048

[vspnat@gmail.com](mailto:vspnat@gmail.com)

### Abstract

A field trial was conducted to determine the studies on different levels and sources of nitrogen on growth and yield of fodder maize (*Zea mays* L.) at Srinivaspur village, Amanikere, Kolar district, Eastern Dry Zone (Zone-5) of Karnataka State, India, during *Kharif* season (July to September - 2021). The experiment was laid out in randomized block design (RBD) with three replications. Among the various treatments evaluated, application of T<sub>10</sub> - 180 Kg N ha<sup>-1</sup> supplied through 50% as urea + 50% as poultry manure + 60:40 Kg of P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O ha<sup>-1</sup> recorded the maximum plant height (247.76 cm), number of leaves 12.06 plant<sup>-1</sup>, leaf length (90.90 cm), leaf width (7 cm), leaf stem ratio (LSR) (0.69) and dry matter production (DMP) (12.38 t ha<sup>-1</sup>), yield characters viz., green fodder yield (GFY) (54.92 t ha<sup>-1</sup>) and green fodder production efficiency (7.12 q ha<sup>-1</sup> day<sup>-1</sup>), and it was on par with T<sub>11</sub> - 180 Kg N supplied through 50% as urea + 50% as sheep manure + 60:40 Kg of P and K ha<sup>-1</sup>. Economics viz., gross returns (Rs. 1,64,762 ha<sup>-1</sup>), net returns (Rs. 1,07,332 ha<sup>-1</sup>) and B: C ratio of 2.86 with (T<sub>10</sub>). The lowest gross returns (Rs. 89,559 ha<sup>-1</sup>), net returns (Rs. 25,387 ha<sup>-1</sup>) and B: C ratio of 1.39 with T<sub>5</sub> - 150 Kg N ha<sup>-1</sup> as sheep manure + 60:40 Kg of P and K ha<sup>-1</sup>. Results indicate that combination of inorganic and organic manure (T<sub>10</sub>) have a significantly greater influence than other treatments.

**Key words :** Nitrogen, poultry manure, sheep manure, growth, yield and economics.

Among the cultivated annual cereal fodder crops, maize is one of the most versatile and multi utility crops having wider adaptability in diverse ecologies. Globally it is known as Queen of cereals because of its highest genetic potential. Maize is the most important fodder crop in the world because of its high yield, high energy forage productivity than that of the other forage crops<sup>13</sup>. Maize being one of the most adaptable emerging crops having wider adaptability under varied agro-climatic conditions has been proved and preferred in terms of green fodder quality and silage making as it provides very palatable, highly succulent, digestible, and nutritionally rich fodder to livestock which is free from anti-metabolites<sup>12</sup>.

It contains 9-10 per cent crude protein, 32.52-33.49 per cent crude fibre, 60-64 per cent neutral detergent fibre, 38-41 per cent acid detergent fibre, 28-30 per cent cellulose and 23-25 per cent hemi cellulose on dry matter basis when harvested at milk to early dough stage. It can be fed as green or dry and makes excellent silage<sup>14</sup>. Fodder maize being a C<sub>4</sub> plant, is very efficient in converting solar energy into dry matter. As a heavy feeder of nutrients, fodder maize productivity is largely dependent on nutrient management. In order to obtain higher crop productivity, management of nutrients through application of organic manure and inorganic fertilizers is an important consideration<sup>6</sup>. The present availability of green fodder is 585 m.t and 494 m.t of dry fodder<sup>2</sup>. The livestock population is increasing @ 1.23 per cent per annum, raising the livestock population approximately to 535.78 million in the country resulting in an increase of 4.6 per cent over the livestock census of

2012. Karnataka is the 9<sup>th</sup> largest state in cattle and buffalo population in the country, accounting for 29 million and it is 4.7 per cent of the total population<sup>9</sup>. In Karnataka, area under cultivation of fodder crops is 3.65 lakh ha while the availability of green fodder is 85 m.t and dry fodder 15 m.t. Demand for green fodder was estimated to be 122 m.t and dry fodder 25.4 m.t per annum<sup>22</sup>.

To meet the fattening livestock, India would require 855 m.t of green fodder, 526 m.t of dry fodder and 56 m.t of feed concentrates. By 2050, the demand for green and dry fodder would be 1012 and 631 m.t, respectively and at the current level of growth in forage resources, there will be 18.4 per cent shortfall deficit in green fodder and 13.2 per cent deficit in dry fodder<sup>2</sup> in the year 2050. The gap in demand and supply may further rise due to consistent growth of livestock population @ 1.23 per cent in the coming years and the demand of green forage supply must rise @ 1.69 per cent per annum to satisfy the deficit; however, the area under cultivated fodder accounts for only 4 per cent of the total cultivated land (8.4 m.ha) in the country, and has remained unchanged over the last few decades<sup>8,11,17</sup>.

Nitrogen is a primary nutrient required by crop plants for their growth and development. It plays a key role in vegetative growth of maize plants. The application of nitrogen not only affects the forage yield of maize, but also improves its quality, mostly its protein content. Application of nitrogen to maize increased growth characters, yield characters and quality characters by increasing nitrogen levels<sup>5,16</sup>. Integrating chemical fertilizer and

organic manure is advantageous in enhancing the availability of nitrogen, phosphorus and potassium in plants, improving soil fertility and productivity on a sustainable basis. These improved soil properties and nutrient availability enhanced the fresh and dry biomass production and improved the crop growth<sup>19</sup>.

A field experiment entitled 'studies on different levels and sources of nitrogen in augmenting the growth and yield of fodder maize (*Zea mays* L.)' was conducted at Srinivaspur village, Amanikere, Kolar district, Eastern Dry Zone (Zone - 5) of Karnataka State, India, during *Kharif* season (July to September - 2021). Geographically this site is situated at 13° 33' N latitude and 78° 21' E longitude with an altitude of +819 m above mean sea level. The soil of experimental field was sandy loam in texture and neutral pH of 7.7. Electrical conductivity was 0.20 dSm<sup>-1</sup> with an organic carbon content of 0.37 %. Available nitrogen was low (110 kg N ha<sup>-1</sup>), phosphorus was medium (20.5 P<sub>2</sub>O<sub>5</sub> ha<sup>-1</sup>) and potassium was medium (202.5 K<sub>2</sub>O ha<sup>-1</sup>). The fodder maize (African Tall) was used as test crop and dibbled by adopting a spacing of 60 × 10 cm. The organic nutrient sources used were poultry manure (PM) and sheep manure (SM) whereas the inorganic sources were urea, single super phosphate (SSP) and muriate of potash (MOP). The experiment was laid out in Randomized Block Design (RBD) with three replications and eleven treatments. Treatments involving a combination of various nutrients from different sources (organic and inorganic) were used which include: T<sub>1</sub> - 120:60:40 kg of NPK ha<sup>-1</sup> (RDF), T<sub>2</sub> - 150:60:40 kg of NPK ha<sup>-1</sup>, T<sub>3</sub> - 180:60:40 kg of NPK ha<sup>-1</sup>, T<sub>4</sub> - 150 kg N ha<sup>-1</sup> as poultry manure +

60:40 kg of P and K ha<sup>-1</sup>, T<sub>5</sub> - 150 kg N ha<sup>-1</sup> as sheep manure + 60:40 kg of P and K ha<sup>-1</sup>, T<sub>6</sub> - 180 kg N ha<sup>-1</sup> as poultry manure + 60:40 kg of P and K ha<sup>-1</sup>, T<sub>7</sub> - 180 kg N ha<sup>-1</sup> as sheep manure + 60:40 kg of P and K ha<sup>-1</sup>, T<sub>8</sub> - 150 kg N ha<sup>-1</sup> supplied through 50% as urea + 50% as poultry manure + 60:40 kg of P and K ha<sup>-1</sup>, T<sub>9</sub> - 150 kg N ha<sup>-1</sup> supplied through 50% as urea + 50% as sheep manure + 60:40 kg of P and K ha<sup>-1</sup>, T<sub>10</sub> - 180 kg N ha<sup>-1</sup> supplied through 50% as urea + 50% as poultry manure + 60:40 kg of P and K ha<sup>-1</sup> and T<sub>11</sub> - 180 kg N ha<sup>-1</sup> supplied through 50% as urea + 50% as sheep manure + 60:40 kg of P and K ha<sup>-1</sup>. The total treatment combinations were 33, each treatment combination was laid out on plot area of 5.4 m × 4 m = 21.6 m<sup>2</sup> and net plot area 4.2 m × 3.8 m.

Observations on growth characters viz., plant height (cm), number of leaves plant<sup>-1</sup>, leaf length (cm), leaf width (cm), leaf stem ratio (LSR) and dry matter production (DMP) (t ha<sup>-1</sup>) were observed from randomly selected five tagged plants in each treatment and the average values of each treatment was calculated and tabulated. Yield characters viz., green fodder yield (GFY) (t ha<sup>-1</sup>) and green fodder production efficiency (q ha<sup>-1</sup> day<sup>-1</sup>). The green fodder yield was computed from the net plot and expressed in t ha<sup>-1</sup>. Green fodder production efficiency was computed as green fodder yield divided by number of days to harvest. Economics viz., cost of cultivation, gross return, net return and benefit cost ratio were also worked out to evaluate the economic benefits of each treatment, based on the existing market prices of inputs and output. The statistical analysis of the field data was

carried out as per the methodology given by Gomez and Gomez<sup>10</sup>. The critical difference were worked out at 5 per cent probability level, wherever the results were significant.

#### *Growth characters :*

The results of the study clearly expressed the influence of various sources and levels of nitrogen on the growth of fodder maize, as transparent from the data on plant height, number of leaves, leaf length, leaf width, leaf stem ratio (LSR) and dry matter production (DMP), and all of the growth parameters at harvest stage of observation in Table-1.

Regarding the different levels of N tried, the remarkable response was noticed up to 180 kg N ha<sup>-1</sup>; the increase in plant height compared to the 120 kg N ha<sup>-1</sup> (T<sub>1</sub>) was 17 per cent, while it was 61.4 and 39.1 per cent for LSR and DMP respectively. N is a major nutrient involved in active cell division and cell elongation, thus enhancing meristematic activity and enhanced uptake of N is likely due to better synthesis of amino acids and protein<sup>22</sup>. The notable rise in growth components and vigorous vegetative growth of fodder maize may be explained by the abundant supply of N, and our finding is well validated by earlier findings of Malik and Paynter<sup>15</sup>.

However, the maximum achievement in growth parameters was registered with INM (T<sub>10</sub> and T<sub>11</sub>) where 50% N of 180 kg ha<sup>-1</sup> was supplied through urea and the remaining N was supplied through poultry manure or sheep manure; the increase in plant height was for a length of 247.76 cm; number

of leaves 12.06 plant<sup>-1</sup>, leaf length (90.90 cm), leaf width (7 cm), 0.69 (LSR) and 12.38 t ha<sup>-1</sup> (DMP). The increase in DMP with INM (T<sub>10</sub>/T<sub>11</sub>) compared with organic manure application alone (sheep and poultry manure) at the same level of 180 kg N ha<sup>-1</sup> (T<sub>6</sub> and T<sub>7</sub>) was 36.91 per cent, while it was 11.79 per cent compared to inorganic N alone @ 180 kg N ha<sup>-1</sup> as urea (T<sub>3</sub>), thus showing the influence and benefit of integrating organic and inorganic in augmenting the growth of fodder maize. It is striking to note that the inorganic source of N alone was better than that of the organic source alone demonstrating the immediate availability of N from inorganic sources, and it is evident from the significant growth parameters with T<sub>10</sub> / T<sub>11</sub>, where part of N was supplied as urea. As N was readily and continuously available from both inorganic and organic sources viz., urea, poultry and sheep manure, the enhanced performance of growth characters in INM treatments is quite understood. Adeniyi and Ojeniyi<sup>1</sup> earlier reported the advantage of INM in fodder maize and endorsed our present findings.

There was a consistent increase in all the growth parameters<sup>21</sup> with the LSR of 0.69. Rasool *et al.*<sup>20</sup> and Nirmal *et al.*,<sup>17</sup> earlier showed the priming effect of organics on the release of N with higher meristematic activity, with a favourable effect on cell division and enlargement. Ayeni and Adetunji<sup>4</sup> also proved that sheep manure provides macro and micronutrients to the root zone, which in turn improves the soil's physical and chemical properties, respectively, in better photosynthetic efficiency. Valadabadi and Farhani<sup>26</sup> showed that N was an integral part of chlorophyll and

enhanced photosynthesis, and it is the fulfilment of our current findings of higher growth characters with treatments having INM at 180 kg N ha<sup>-1</sup>. So many workers reported the advantage of harmonizing organics and inorganics in enhancing the various growth characteristics of fodder maize<sup>7</sup> (on plant height; Sharma *et al.*,<sup>21</sup> on LSR; Asif Iqbal *et al.*,<sup>3</sup> on DMP) and thus contribute confidence to our present findings.

#### *Yield characters :*

As the yield of any crop is normally the manifestation of all growth components, the same is true, with our present finding also Table 2. The highest green fodder production efficiency of 7.12 q ha<sup>-1</sup> day<sup>-1</sup> was noticed with T<sub>10</sub> however, on par with T<sub>11</sub>, which was higher

by 42 per cent compared with T<sub>1</sub> (120 kg N ha<sup>-1</sup>). Similarly, T<sub>10</sub> / T<sub>11</sub> achieved the highest green fodder yield (54.92 t ha<sup>-1</sup>) which was higher by 11.7 per cent compared to T<sub>3</sub> indicating the effectiveness of integration and response to higher levels of N. Likewise, at 150 kg N ha<sup>-1</sup> (T<sub>2</sub>), the increase with T<sub>8</sub> / T<sub>9</sub> (150 kg N ha<sup>-1</sup> supplied through 50% as urea + 50% as sheep manure + 60:40 kg of P and K ha<sup>-1</sup>) over T<sub>2</sub> was 10.5 per cent, again demonstrating the advantage of an organic source integrated with an inorganic source of N at any level. An increase of 23.2 per cent green fodder yield was noticed with T<sub>2</sub> over T<sub>1</sub> and 18.8 per cent with T<sub>3</sub> over T<sub>2</sub>, thus indicating the response to incremental N. As observed in our present study, a similar growth and yield pattern was earlier reported by many authors<sup>1,26</sup>. As N is a constituent of chlorophyll

Table-1. Effect of different levels and sources of N on growth characters at harvest stage of fodder maize

Treatments	Plant height (cm)	Number of leaves (plant <sup>-1</sup> )	Leaf Length (cm)	Leaf width (cm)	LSR	DMP (t ha <sup>-1</sup> )
T <sub>1</sub>	197.26	10.30	80.66	5.80	0.22	6.65
T <sub>2</sub>	217.16	10.96	84.73	6.30	0.37	8.81
T <sub>3</sub>	237.06	11.73	88.76	6.73	0.57	10.92
T <sub>4</sub>	196.70	10.20	79.96	5.80	0.21	6.71
T <sub>5</sub>	196.53	10.13	79.86	5.80	0.20	6.84
T <sub>6</sub>	207.26	10.63	82.83	6.0	0.31	7.81
T <sub>7</sub>	207.10	10.60	82.76	6.0	0.30	7.70
T <sub>8</sub>	227.30	11.43	87.00	6.53	0.49	9.87
T <sub>9</sub>	227.16	11.33	86.96	6.50	0.48	9.80
T <sub>10</sub>	247.76	12.06	90.90	7.0	0.69	12.38
T <sub>11</sub>	247.26	12.03	90.86	6.96	0.68	12.29
S.Em±	3.29	0.09	0.58	0.04	0.01	0.29
CD at 5%	9.72	0.27	1.73	0.14	0.05	0.85

Table-2. Effect of different levels and sources of N on yield characters and economics of fodder maize

Treatments	GFY (t ha <sup>-1</sup> )	Green fodder production efficiency (q ha <sup>-1</sup> day <sup>-1</sup> )	Cost of cultivation (Rs.ha <sup>-1</sup> )	Gross returns (Rs.ha <sup>-1</sup> )	Net returns (Rs.ha <sup>-1</sup> )	B : C ratio
T <sub>1</sub>	30.20	4.02	50196	90628	40432	1.80
T <sub>2</sub>	39.33	5.27	50595	118008	67412	2.33
T <sub>3</sub>	48.46	6.44	50993	145384	94390	2.85
T <sub>4</sub>	30.08	4.02	61323	90255	28932	1.47
T <sub>5</sub>	29.85	3.95	64172	89559	25387	1.39
T <sub>6</sub>	34.77	4.65	63866	104329	40462	1.63
T <sub>7</sub>	34.66	4.60	67529	104000	36470	1.54
T <sub>8</sub>	44.13	5.77	55959	132413	76453	2.36
T <sub>9</sub>	43.98	5.75	57475	131958	74482	2.29
T <sub>10</sub>	54.92	7.12	57430	164762	107332	2.86
T <sub>11</sub>	54.91	7.03	59261	164738	105476	2.77
S.Em±	1.47	0.15	-	-	-	-
CD at 5%	4.36	0.46	-	-	-	-

•Green fodder cost Rupees 3/- per kg

and an essential functionary of the growth of plants, its continuous availability during the entire duration of the crop results in higher leaf and stem growth<sup>24</sup>. In our present study, N was continuously available from an inorganic and organic source (vide Table-2), and hence a significant increase in T<sub>10</sub> and T<sub>11</sub> was noticed. Similar findings of the response of fodder maize were earlier demonstrated by Asif Iqbal *et al.*<sup>3</sup> and Yadav *et al.*<sup>27</sup>.

#### Economics :

The economics for various treatments was drawn using the cost involved and market

price of green fodder showed in Table-2. The total cost of cultivation varies between Rs. 59, 261 ha<sup>-1</sup> to Rs. 50, 196 ha<sup>-1</sup>. Among the different treatments, INM plot recorded the highest green fodder yield at 54.92 t ha<sup>-1</sup> (T<sub>10</sub>) registered highest gross return of Rs. 1, 64, 762 ha<sup>-1</sup>, net return of Rs. 1, 07, 332 ha<sup>-1</sup> and B: C ratio of 2.86. This was followed by T<sub>11</sub> and in turn by T<sub>3</sub> which gave a gross return of Rs. 1, 64, 738 and 1, 45, 384 ha<sup>-1</sup> respectively. The lowest gross returns (Rs. 89, 559 ha<sup>-1</sup>) were registered with T<sub>5</sub>. Net return from different treatments followed a similar trend as that of gross return. The highest net return

of Rs. 1, 07, 332 ha<sup>-1</sup> was obtained due to application of T<sub>10</sub>. This was followed by T<sub>11</sub> and T<sub>3</sub> which gave a net return of Rs. 1, 05, 476 and 94, 390 ha<sup>-1</sup> respectively. The lowest net return of Rs. 25, 387 and 28, 932 ha<sup>-1</sup> was registered with T<sub>5</sub> and T<sub>4</sub>, respectively.

From the results obtained in the experiment conducted in the Eastern dry zone of Srinivasapur, Karnataka State, it may be concluded that fodder maize variety African Tall responded well up to 180 kg N ha<sup>-1</sup> applied as 50% through urea and 50% through either poultry manure or sheep manure along with 60 kg P<sub>2</sub>O<sub>5</sub> and 40 kg K<sub>2</sub>O as SSP and MOP respectively that gave the maximum green fodder yield of 54.92 t ha<sup>-1</sup> and net returns of Rs. 1, 07, 332 ha<sup>-1</sup> and B: C ratio of 2.86. (T<sub>10</sub>) showed best performance than all the other treatments in increasing the growth and yield characters of fodder maize.

#### References :

1. Adeniyi, O.N. and S.O. Ojeniyi (2003). *Journal of Agricultural Research*, 4: 191-197.
2. Anonymous, (2013). Indian grassland and fodder research institute. 251.
3. Asif Iqbal, Muhammad Aamir Iqbal, Ali Raza, Nadeem Akbar, Rana Nadeem Abbas and Haroon Zaman Khan (2014). *American-Eurasian. Journal of agricultural and environmental sciences*, 14(8): 744-747.
4. Ayeni, L.S. and M.T. Adetunji (2010). *Journal of Natural Sciences.*, 8(1): 60-67.
5. Aziz Khan, Fazal Munsif, Kashif Akhtar, Muhammad Zahir Afridi, Zahoor, Zahoor Ahmad, Shah Fahad, Rizwan Ullah, Faheem Ahmed Khan, Mairaj Din (2014). *American Journal of Plant Science* 5: 2323-2329.
6. Biswasi, S.K., A.K. Barik, D.K. Bastia, B. Dalei, L. Nayak and M. Ray (2020). *International Journal Bio-resources and Stress Management*. 11(5): 465-471.
7. Cheema, M.A., W. Farhad, M.F. Saleem, H.Z. Khan, A. Munir, M.A. Wahid, F. Rasul and H.M. Hammad (2010). *Crop & Environment*, 1(1): 49-52.
8. Dagar, J.C., P.K. Ghosh, S.K. Mohanta, J.B. Singh, D. Vijay Kumar, V.K. Yadav and S. Kumar (Eds.). (2017). Potentials for fodder production in degraded lands. Approaches towards fodder security in India. 333-364. Studera Press New Delhi.
9. GOI, (2019). Department of Animal Husbandry and Dairying, Ministry of Fisheries, Animal Husbandry and Dairying.
10. Gomez, A.K and A.A. Gomez (1984). Statistical procedure for agricultural research. International Rice Research Institute, Manila, Phillippines. pp. 304-305.
11. Halli, H.M., S.S. Rathore, N. Manjunatha and V. K. Wasnik (2018). *International Journal of Current Microbiology and Applied Science*, 7(02): 1912-1921.
12. Kumar, Naveen, Satpal, Suresh Kumar, Uma Devi, J.M. Sutaliya and Shweta (2020). *Forage Research*. 46: 10-21.
13. Kumar, S., R.K. Arya, N. Singh and Satpal (2021). *Forage Research*. 47: 125-129.
14. Kumar Sunil, R.K. Agrawal, A.K. Dixit, K. Rai, Arvind and S.K. Rai (2012). Forage crops and their management. Indian grassland and fodder research institute, Jhansi-Uttar Pradesh, India. 60.
15. Malik, R.K. and B. Paynter (2010). *Soil Solution for Acquiring World*, 1(6):187-

- 189.
16. Manjunatha, S.B., V.V. Angadi, Y.B. Palled, S.V. Hosamani, H.B. Babalad, M.H. Hosamani, D.H. Mudakavi and H.T. Channal (2017). *Karnataka Journal Agricultural Science* 26(4): 511-513.
17. Meena, L.R., S.A. Kochewad, V. Kumar, S. Malik, S. Kumar, L.K. Meena, A.L. Meena and A.S. Panwar (2018). *Range Management & Agroforestry*, 39(2): 313-318.
18. Nirmal, S.S., D.D. Dudhade, A.V. Solanke, S.R. Gadakh, B.D. Bhakare, R.R. Hasure and S.B. Gore (2016). *International Journal of Science Environment and Technology*, 5(5): 2999-3004.
19. Rajashekhara Rao, B.K., K.L. Sahrawat, S.P. Wani and G. Pardhasaradhy (2010). *International Journal of Soil Science*, 5(4): 216-225.
20. Rasool, S., R.H. Kanth, S. Hamid, W. Raja, B.A. Alie and Z.A. Dar (2015). *American Journal of Experimental Agriculture*. 13(2): 1-9.
21. Sharma, P.K., V.P. Kalra and U.S. Tiwana (2016). *Agricultural Research Journal*. 53(3): 355-359.
22. Shekara, B.G., P. Mahadevu, N.M. Chikkarugi and N. Manasa (2019). Fodder and livestock scenario in Karnataka. 79-90.
23. Singh, Bahadur, M.K. Kaushik and H.K. Sumeriya (2015). *Annals of Agri-Bio Research* 20: 30-32.
24. Subrahmanya, D.J., Rakesh Kumar, P.S. Pyati, Hardev Ram, R.K. Meena and Akanksha Tamta (2019). *Forage Research*. 45(2): 127-131.
25. Valadabadi, S.A. and H.A. Farhani (2010). *Journal of Agricultural Extension and Rural Development*. 2(3): 040-047.
26. Vasanthi, D. and K. Kumaraswamy (2000). *Journal of Indian Society of Soil Science* 48: 510-515.
27. Yadav Sunitha, Ramawatar Meena, Seema, Sandeep Kumar and D.K. Sharma (2019). *Indian Journal Agricultural Science*. 89(11): 227-229.